



A unified analysis of tangent sets to determinantal varieties

Yan Yang

SIAM OP26, Edinburgh

June 2, 2026

State Key Laboratory of Mathematical Sciences
Academy of Mathematics and Systems Science
Chinese Academy of Sciences, China

Joint work with **Dr. Bin Gao** and **Prof. Ya-xiang Yuan**

♥ Registration and travel support for this presentation was provided by
the **Society for Industrial and Applied Mathematics (SIAM)**
and **Academy of Mathematics and Systems Science (AMSS)**

siam
Society for Industrial and Applied Mathematics



Academy of Mathematics and Systems Science
Chinese Academy of Sciences

- 1 Background: low-rank optimization
- 2 Variational analysis of low-rank sets
- 3 First- and second-order tangent sets
- 4 Tangent sets bridge optimization landscapes
- 5 Second-order optimality on bounded-rank matrices

Background: low-rank optimization

Optimization on the fixed-rank manifold $r < \min\{m, n\}$

$$\min f(X) \quad \text{s.t. } X \in \mathcal{M}_r := \{X \in \mathbb{R}^{m \times n} \mid \text{rank}(X) = r\}$$

- $\mathcal{M}_r$ is a smooth manifold [Helmke-Shayman'95]
- Riemannian optimization algorithms [Vandereycken'13]
- $\mathcal{M}_r$ is **not closed**

Optimization on the fixed-rank manifold $r < \min\{m, n\}$

$$\min f(X) \quad \text{s.t. } X \in \mathcal{M}_r := \{X \in \mathbb{R}^{m \times n} \mid \text{rank}(X) = r\}$$

- $\mathcal{M}_r$ is a smooth manifold [Helmke-Shayman'95]
- Riemannian optimization algorithms [Vandereycken'13]
- $\mathcal{M}_r$ is **not closed**

Optimization on the set of bounded-rank matrices $r < \min\{m, n\}$

$$\min f(X) \quad \text{s.t. } X \in \mathcal{M}_{\leq r} := \{X \in \mathbb{R}^{m \times n} \mid \text{rank}(X) \leq r\}$$

- $\mathcal{M}_{\leq r}$ is **closed**
- $\mathcal{M}_{\leq r}$ is a real-algebraic variety: **determinantal variety** [Harris'92]
- Finding first-order stationary points [Schneider-USchmajew'15; Levin-Kileel-Boumal'22; '24; Olikier-Absil'22; '23; '24]

Optimization on the fixed-rank manifold $r < \min\{m, n\}$

$$\min f(X) \quad \text{s.t. } X \in \mathcal{M}_r := \{X \in \mathbb{R}^{m \times n} \mid \text{rank}(X) = r\}$$

- $\mathcal{M}_r$ is a smooth manifold [Helmke-Shayman'95]
- Riemannian optimization algorithms [Vandereycken'13]
- $\mathcal{M}_r$ is **not closed**

Optimization on the set of bounded-rank matrices $r < \min\{m, n\}$

$$\min f(X) \quad \text{s.t. } X \in \mathcal{M}_{\leq r} := \{X \in \mathbb{R}^{m \times n} \mid \text{rank}(X) \leq r\}$$

- $\mathcal{M}_{\leq r}$ is **closed**
- $\mathcal{M}_{\leq r}$ is a real-algebraic variety: **determinantal variety** [Harris'92]
- Finding first-order stationary points [Schneider-USchmajew'15; Levin-Kileel-Boumal'22; '24; Olikier-Absil'22; '23; '24]

A unified variational analysis for low-rank sets ?

$$\begin{array}{ll} \min & f(X) \\ \text{s.t.} & X \in \mathcal{M}_{\leq r} \end{array}$$

- Recommendation system: movie ratings [Frolov-Oseledets'17]
- Hyperspectral Images [Zhang-He-Zhang-Shen-Yuan'13; Zhuang-Fu-Ng'21]
- Image and video inpainting [Bertalmio-Sapiro-Caselles-Ballester'00; Fu-Ruan-Luo-An-Jin'21; Luo-Zhao-Li-Ng-Meng'23]
- EEG (brain signals) data [Mørup-Hansen-Herrmann-Parnas-Arnfred'06; Kong-Kong-Fan-Zhao-Cichoki'17]
- Magnetic resonance imaging (MRI) [Banco-Aeron-Hoge'16; Choi-Bao-Zhang'18; Fessler'20]
- Data analysis, e.g., weather forecast [Loucheur-Absil-Journée'23] and exoplanet detection [Daglayan-Vary-Absil-Cantalloube-Christiaens-Gillis-Jacques-Leplat-Absil'24]

Bounded-rank matrices + an additional constraint

$$\begin{array}{ll} \min & f(X) \\ \text{s.t.} & X \in \mathcal{M}_{\leq r} \cap \mathcal{H} \end{array}$$


Bounded-rank matrices + an additional constraint

$$\begin{array}{ll} \min & f(X) \\ \text{s.t.} & X \in \mathcal{M}_{\leq r} \cap \mathcal{H} \end{array}$$

Low rank + oblique manifold

- Low-rank data fitting on sphere [Chu-Del Buono-Lopez-Politi'05 SIMAX]

$$\mathcal{H} = \text{Ob}(m, n) := \{X \in \mathbb{R}^{m \times n} \mid \text{diag}(XX^\top) - \mathbf{1} = \mathbf{0}\}$$



Low rank + Frobenius sphere

- Approximation of graph similarity matrices [Cason-Absil-Van Dooren'13 LAA]

$$\mathcal{H} = S_F(m, n) := \{X \in \mathbb{R}^{m \times n} \mid \|X\|_F^2 - 1 = 0\}$$

Low rank + hyperbolic manifold

- Approximation of hyperbolic embeddings [Jawanpuria-Meghwanshi-Mishra'19 CDC]

$$\mathcal{H} = \mathbb{H}_{n-1}^m := \{X \in \mathbb{R}^{m \times n} \mid \text{diag}(X \text{Diag}(-1, 1, \dots, 1)X^\top) - \mathbf{1} = \mathbf{0}\}$$

Low rank + additional constraints (cont'd)

Low rank + Frobenius sphere

- Approximation of graph similarity matrices [Cason-Absil-Van Dooren'13 LAA]

$$\mathcal{H} = \mathcal{S}_F(m, n) := \{X \in \mathbb{R}^{m \times n} \mid \|X\|_F^2 - 1 = 0\}$$

Low rank + hyperbolic manifold

- Approximation of hyperbolic embeddings [Jawanpuria-Meghwanshi-Mishra'19 CDC]

$$\mathcal{H} = \mathbb{H}_{n-1}^m := \{X \in \mathbb{R}^{m \times n} \mid \text{diag}(X \text{Diag}(-1, 1, \dots, 1)X^\top) - \mathbf{1} = \mathbf{0}\}$$

Low rank + SDPs

- Low-rank semidefinite programming [Journée-Bach-Absil-Sepulchre'10 SIOPT]

$$\min \quad \langle C, X \rangle$$

$$\text{s.t.} \quad X \in \mathcal{M}_{\leq r} \cap \mathcal{S}^+(n) \cap \mathcal{U}$$

$$\text{where } \mathcal{U} = \{X \in \mathbb{R}^{n \times n} \mid \mathcal{A}(X) = b\}$$

Low rank + additional constraints (cont'd)

Low rank + Frobenius sphere

- Approximation of graph similarity matrices [Cason-Absil-Van Dooren'13 LAA]

$$\mathcal{H} = \mathcal{S}_F(m, n) := \{X \in \mathbb{R}^{m \times n} \mid \|X\|_F^2 - 1 = 0\}$$

Low rank + hyperbolic manifold

- Approximation of hyperbolic embeddings [Jawanpuria-Meghwanshi-Mishra'19 CDC]

$$\mathcal{H} = \mathbb{H}_{n-1}^m := \{X \in \mathbb{R}^{m \times n} \mid \text{diag}(X \text{Diag}(-1, 1, \dots, 1)X^\top) - \mathbf{1} = \mathbf{0}\}$$

Low rank + SDPs

- Low-rank semidefinite programming [Journée-Bach-Absil-Sepulchre'10 SIOPT]

$$\min \quad \langle C, X \rangle$$

$$\text{s.t.} \quad X \in \mathcal{M}_{\leq r} \cap \mathcal{S}^+(n) \cap \mathcal{U}$$

$$\text{where } \mathcal{U} = \{X \in \mathbb{R}^{n \times n} \mid \mathcal{A}(X) = b\}$$

☆ First- and second-order optimality conditions

Variational analysis of low-rank sets

Constraint-coupled optimization

$$\begin{array}{ll} \min & f(X) \\ \text{s. t.} & X \in \mathcal{M}_{\leq r} \cap \mathcal{H} \end{array}$$

Challenges

- Inherent **non-smoothness** of $\mathcal{M}_{\leq r}$ [Olikier-Absil'22 SVVA; Levin-Kileel-Boumal'23; '25 MP]
- **Geometry** of the coupled region $\mathcal{M}_{\leq r} \cap \mathcal{H}$
 - **Complicated** Bouligand-tangent cone
 - **Unknown** second-order tangent set

Tangent and normal cones to $\mathcal{M}_{\leq r}$

- Mordukhovich normal cone [Luke'13]
- Bouligand tangent cone and Frèchet normal cone [Cason-Absil-Van Dooren'13]
- Clarke tangent cone and Clarke normal cone [Hosseini-Uschmajew'19; Li-Song-Xiu'19]
- Continuity of tangent cones [Olikier-Absil'22]

Tangent and normal cones to $\mathcal{M}_{\leq r}$

- Mordukhovich normal cone [Luke'13]
- Bouligand tangent cone and Frèchet normal cone [Cason-Absil-Van Dooren'13]
- Clarke tangent cone and Clarke normal cone [Hosseini-Uschmajew'19; Li-Song-Xiu'19]
- Continuity of tangent cones [Olikier-Absil'22]

First-order geometry of $\mathcal{M}_{\leq r} \cap \mathcal{H}$

- Mordukhovich normal cone: $\mathcal{H} = \text{Sym}(n)$ [Tam'17]
- Frèchet normal cone: $\mathcal{H} = \text{Sym}(n) \cap \mathcal{B}_i$ [Li-Xiu-Zhou'20 JOTA]
 $\mathcal{B}_1 = \{X \mid \|X\|_{\text{F}}^2 \leq 1\}, \mathcal{B}_2 = \{X \mid -tI_n \preceq X \preceq tI_n\}, \mathcal{B}_3 = \{X \mid X \succeq 0, \text{tr}(X) = 1\}$
- Bouligand tangent cone: $\mathcal{H} = \mathcal{S}^+(n)$ [Levin-Kileel-Boumal'25 MP]

Tangent and normal cones to $\mathcal{M}_{\leq r}$

- Mordukhovich normal cone [Luke'13]
- Bouligand tangent cone and Frèchet normal cone [Cason-Absil-Van Dooren'13]
- Clarke tangent cone and Clarke normal cone [Hosseini-Uschmajew'19; Li-Song-Xiu'19]
- Continuity of tangent cones [Olikier-Absil'22]

First-order geometry of $\mathcal{M}_{\leq r} \cap \mathcal{H}$

- Mordukhovich normal cone: $\mathcal{H} = \text{Sym}(n)$ [Tam'17]
- Frèchet normal cone: $\mathcal{H} = \text{Sym}(n) \cap \mathcal{B}_i$ [Li-Xiu-Zhou'20 JOTA]

$$\mathcal{B}_1 = \{X \mid \|X\|_F^2 \leq 1\}, \mathcal{B}_2 = \{X \mid -tI_n \preceq X \preceq tI_n\}, \mathcal{B}_3 = \{X \mid X \succeq 0, \text{tr}(X) = 1\}$$

- Bouligand tangent cone: $\mathcal{H} = \mathcal{S}^+(n)$ [Levin-Kileel-Boumal'25 MP]
- Bouligand tangent cone: $\mathcal{H} = \{X \in \mathbb{R}^{m \times n} \mid \|X\|_F^2 = 1\}$ [Cason-Absil-Van Dooren'13 LAA]
- Frèchet normal cone: $\mathcal{H} = \{X \in \mathbb{R}^{m \times n} \mid \mathcal{A}(X) = b\}$ [Li-Luo'23 SIOPT]

★ Bouligand tangent cone and Frèchet normal cone:

$$\mathcal{H} = \{X \in \mathbb{R}^{m \times n} \mid h(X) = 0\} \text{ with orthogonally invariant } h \text{ [Yang-Gao-Yuan'26 MP]}$$

Tangent and normal cones to $\mathcal{M}_{\leq r}$

- Mordukhovich normal cone [Luke'13]
- Bouligand tangent cone and Frèchet normal cone [Cason-Absil-Van Dooren'13]
- Clarke tangent cone and Clarke normal cone [Hosseini-Uschmajew'19; Li-Song-Xiu'19]
- Continuity of tangent cones [Olikier-Absil'22]

First-order geometry of $\mathcal{M}_{\leq r} \cap \mathcal{H}$

- Mordukhovich normal cone: $\mathcal{H} = \text{Sym}(n)$ [Tam'17]
- Frèchet normal cone: $\mathcal{H} = \text{Sym}(n) \cap \mathcal{B}_i$ [Li-Xiu-Zhou'20 JOTA]

$$\mathcal{B}_1 = \{X \mid \|X\|_{\text{F}}^2 \leq 1\}, \mathcal{B}_2 = \{X \mid -tI_n \preceq X \preceq tI_n\}, \mathcal{B}_3 = \{X \mid X \succeq 0, \text{tr}(X) = 1\}$$


- Bouligand tangent cone: $\mathcal{H} = \mathcal{S}^+(n)$ [Levin-Kileel-Boumal'25 MP]
- Bouligand tangent cone: $\mathcal{H} = \{X \in \mathbb{R}^{m \times n} \mid \|X\|_{\text{F}}^2 = 1\}$ [Cason-Absil-Van Dooren'13 LAA]
- Frèchet normal cone: $\mathcal{H} = \{X \in \mathbb{R}^{m \times n} \mid \mathcal{A}(X) = b\}$ [Li-Luo'23 SIOPT]

★ Bouligand tangent cone and Frèchet normal cone:

$$\mathcal{H} = \{X \in \mathbb{R}^{m \times n} \mid h(X) = 0\} \text{ with orthogonally invariant } h \text{ [Yang-Gao-Yuan'26 MP]}$$

A unified analysis remains unclear

Tangent sets and optimization



A unified analysis of tangent sets

Set	Format	First-order	Second-order
$\mathcal{M}$	Assumption 1	Theorem 1	Theorem 1
$\mathcal{M}_{\leq r}$	matrix	[Schneider-USchmajew'15 SIOPT]	Proposition 1
$\mathcal{M}_{\leq r}^{\text{ht}}$	hierarchical Tucker	Proposition 2	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tc}}$	Tucker	[Gao-Peng-Yuan'25 MP]	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tt}}$	tensor train	[Kutschan'18 LAA]	Proposition 2
$\mathcal{S}_{\leq r}(n)$	symmetric matrix	[Li-Xiu-Zhou'20 JOTA]	Proposition 3
$\mathcal{S}_{\leq r}^+(n)$	PSD matrix	[Levin-Kileel-Boumal'25 MP]	Proposition 4
Intersection of sets	Structured set	First-order	Second-order
$\mathcal{M} \cap \mathcal{K}$	Assumption 2	Theorem 2	Theorem 2
$\mathcal{M}_{\leq r} \cap \mathcal{H}$	$\mathcal{H}$ is an affine manifold	[Li-Luo'23 SIOPT]	Appendix C.1
$\mathcal{M}_{\leq r} \cap \mathcal{H}$	$\mathcal{H}$ is orthogonally invariant	[Yang-Gao-Yuan'25]	Appendix C.2
$\mathcal{M}_{\leq r} \cap \mathcal{H}$	$\mathcal{H}$ is hyperbolic	Appendix C.3	Appendix C.3
$\mathcal{S}_{\leq r}(n) \cap \mathcal{U}$	$\mathcal{U} = \{X \mid \ X\ _{\text{F}}^2 = 1\}$	Appendix D.2	Appendix D.2
$\mathcal{S}_{\leq r}^+(n) \cap \mathcal{U}$	$\mathcal{U} = \{X \mid \mathcal{A}(X) = b\}$	[Levin-Kileel-Boumal'25 MP]	Appendix D.3

Bouligand tangent cone

$$T_{\mathcal{X}}(X) = \{\eta \in \mathcal{E} : \exists t_i \rightarrow 0, \text{ s. t. } \text{dist}(X + t_i \eta, \mathcal{X}) = o(t_i)\}$$

Fréchet and Mordukhovich normal cones

$$T_{\mathcal{X}}(X) \xrightarrow{\text{polar}} \hat{N}_{\mathcal{X}}(X) \xrightarrow{\lim} N_{\mathcal{X}}(X)$$

Second-order tangent set in the direction η

$$T_{\mathcal{X}}^2(X, \eta) = \left\{ \zeta \in \mathcal{E} : \exists t_i \rightarrow 0, \text{ s. t. } \text{dist}\left(X + t_i \eta + \frac{1}{2} t_i^2 \zeta, \mathcal{X}\right) = o(t_i^2) \right\}$$

Directional derivative in the direction η

$$h'(X; \eta) := \lim_{t \rightarrow 0} \frac{h(X + t\eta) - h(X)}{t}$$

Parabolic second-order directional derivative in the direction (η, ζ)

$$h''(X; \eta, \zeta) := \lim_{t \rightarrow 0} \frac{h(X + t\eta + \frac{1}{2} t^2 \zeta) - h(X) - th'(X; \eta)}{\frac{1}{2} t^2}$$

First-order geometry of low-rank sets

$\mathcal{M}_s$ as a smooth manifold based on the SVD $X = U\Sigma V^\top$

$$\begin{aligned}T_{\mathcal{M}_s}(X) &= \left\{ [U \ U_\perp] \begin{bmatrix} \mathbb{R}^{s \times s} & \mathbb{R}^{s \times (n-s)} \\ \mathbb{R}^{(m-s) \times s} & \mathbb{0}^{(m-s) \times (n-s)} \end{bmatrix} [V \ V_\perp]^\top \right\} \\N_{\mathcal{M}_s}(X) &= \left\{ [U \ U_\perp] \begin{bmatrix} \mathbb{0}^{s \times s} & \mathbb{0}^{s \times (n-s)} \\ \mathbb{0}^{(m-s) \times s} & \mathbb{R}^{(m-s) \times (n-s)} \end{bmatrix} [V \ V_\perp]^\top \right\}\end{aligned}$$

Geometry of $\mathcal{M}_{\leq r}$ at $X = U\Sigma V^\top \in \mathcal{M}_s$ [Luke'13; Cason-Absil-Van Dooren'13; Schneider-Uschmajew'15]

$$T_{\mathcal{M}_{\leq r}}(X) = T_{\mathcal{M}_s}(X) + \{R \in N_X \mathcal{M}_s : \text{rank}(R) \leq r - s\}$$

$$\hat{N}_{\mathcal{M}_{\leq r}}(X) = \begin{cases} N_{\mathcal{M}_s}(X), & \text{if } s = r \\ \{0\}, & \text{if } s < r \end{cases}$$

$$N_{\mathcal{M}_{\leq r}}(X) = \{Y \in N_{\mathcal{M}_s}(X) : \text{rank}(Y) \leq \min\{m, n\} - r\}$$

All the cone mappings are not continuous at **singular** points [Olikier-Absil'22]


First- and second-order tangent sets

First- and second-order tangent sets

$$\mathcal{T}_{\mathcal{X}}(X) = \{\eta \in \mathcal{E} : \exists t_i \rightarrow 0, \text{ s. t. } \text{dist}(X + t_i\eta, \mathcal{X}) = o(t_i)\}$$

$$\mathcal{T}_{\mathcal{X}}^2(X, \eta) = \left\{ \zeta \in \mathcal{E} : \exists t_i \rightarrow 0, \text{ s. t. } \text{dist}\left(X + t_i\eta + \frac{1}{2}t_i^2\zeta, \mathcal{X}\right) = o(t_i^2) \right\}$$

Approximations to the set $\mathcal{X} = \{(x, y) \in \mathbb{R}^2 \mid y = x^2 + x^3\}$



Challenge when $\mathcal{X} = \mathcal{M}_{\leq r} = \{X \in \mathbb{R}^{m \times n} \mid \text{rank}(X) \leq r\}$

- The geometry of $\mathcal{M}_{\leq r}$ is **complicated** [Olikier-Absil'22]
- The mapping $\text{rank}(\cdot)$ is **not continuous**

A new perspective

$$\mathcal{M}_{\leq r} = \{X \in \mathbb{R}^{m \times n} \mid \sigma_{r+1}(X) = 0\}$$

☹ σ_{r+1} is both non-differentiable and non-convex

😊 σ_{r+1} is Lipschitz continuous [Weyl 1912]

$$|\sigma_{r+1}(X) - \sigma_{r+1}(X + \Delta)| \leq \|\Delta\|_2$$

😊 σ_{r+1} controls the distance to $\mathcal{M}_{\leq r}$

$$\text{dist}(\tilde{X}, \mathcal{M}_{\leq r}) \leq (\min\{m, n\} - r)^{1/2} \sigma_{r+1}(\tilde{X})$$

Error bound condition at X

Let $\mathcal{M} := \{\tilde{X} \in \mathbb{R}^q \mid c_1(\tilde{X}) = 0, c_2(\tilde{X}) \leq 0\}$. There exists a neighborhood $\mathcal{B}$ of $X \in \mathcal{M}$ and a constant $\rho > 0$ such that

$$\text{dist}(\tilde{X}, \mathcal{M}) \leq \rho \|(c_1(\tilde{X}), [c_2(\tilde{X})]_+)\|_2, \text{ for all } \tilde{X} \in \mathcal{B}$$

Theorem

Let $\mathcal{M} = \{\tilde{X} \in \mathbb{R}^q \mid c_1(\tilde{X}) = 0, c_2(\tilde{X}) \leq 0\}$ with c_1 and c_2 satisfying

- error bound condition
- local Lipschitz property

Define the active index set $I_0(X) := \{j \in \{1, \dots, n_2\} \mid c_2(X)_j = 0\}$

(i) (First-order) If c_1 and c_2 are directionally differentiable at X , then

$$\mathbf{T}_{\mathcal{M}}(X) = \{\eta \in \mathbb{R}^q \mid c'_1(X; \eta) = 0, c'_2(X; \eta)_j \leq 0 \text{ for all } j \in I_0(X)\}$$

(ii) (Second-order) If, in addition, c_1 and c_2 admit parabolic second-order directional derivatives at X , then for any $\eta \in \mathbf{T}_{\mathcal{M}}(X)$

$$\mathbf{T}_{\mathcal{M}}^2(X; \eta) = \left\{ \zeta \in \mathbb{R}^q \mid c''_1(X; \eta, \zeta) = 0, c''_2(X; \eta, \zeta)_j \leq 0 \text{ for all } j \in I_1(X; \eta) \right\}$$

where $I_1(X; \eta) := \{j \in I_0(X) \mid c'_2(X; \eta)_j = 0\}$

Low-rank matrices

$$\mathcal{M}_{\leq r} = \{X \in \mathbb{R}^{m \times n} \mid \sigma_{r+1}(X) = 0\}$$

- $\|Y - \mathcal{P}_{\mathcal{M}_{\leq r}}(Y)\|_F = \sqrt{\sum_{i=r+1}^{\min\{m,n\}} \sigma_i^2} \leq (\min\{m, n\} - r)^{1/2} \sigma_{r+1}$
- $\mathcal{T}_{\mathcal{M}_{\leq r}}(X) = \{\eta \in \mathbb{R}^{m \times n} \mid \sigma'_{r+1}(X; \eta) = 0\}$

Low-rank matrices

$$\mathcal{M}_{\leq r} = \{X \in \mathbb{R}^{m \times n} \mid \sigma_{r+1}(X) = 0\}$$

- $\|Y - \mathcal{P}_{\mathcal{M}_{\leq r}}(Y)\|_F = \sqrt{\sum_{i=r+1}^{\min\{m,n\}} \sigma_i^2} \leq (\min\{m, n\} - r)^{1/2} \sigma_{r+1}$
- $\mathcal{T}_{\mathcal{M}_{\leq r}}(X) = \{\eta \in \mathbb{R}^{m \times n} \mid \sigma'_{r+1}(X; \eta) = 0\}$

Low-rank tensors hierarchical Tucker, Tucker, and tensor train

$$\mathcal{M}_{\leq \mathbf{r}}^{\text{ht}} = \bigcap_{t \in T} \text{ten}_{\leq r_t}^{\text{ht}} \left(\mathbb{R}_{\leq r_t}^{n_t \times n_{t-}} \right) = \left\{ \mathbf{X} \in \mathbb{R}^{n_1 \times n_2 \times \cdots \times n_d} \mid \sigma_{r_t+1}(X_{(t)}^{\text{ht}}) = 0 \text{ for } t \in T \right\}$$

- $\|\mathbf{Y} - \mathcal{P}_{\leq \mathbf{r}}^{\text{HOSVD}}(\mathbf{Y})\|_F \leq \rho' \sqrt{\sum_{t \in T} \sigma_{r_t+1}^2(Y_{(t)}^{\text{ht}})}$
- $\mathcal{T}_{\mathcal{M}_{\leq \mathbf{r}}^{\text{ht}}}(\mathbf{X}) = \bigcap_{t \in T} \text{ten}_{(t)}^{\text{ht}} \left(\mathcal{T}_{\mathcal{R}_t}(X_{(t)}^{\text{ht}}) \right)$ with $\mathcal{R}_t := \mathbb{R}_{\leq r_t}^{n_t \times n_{t-}}$

Low-rank **symmetric matrices**

$$\mathcal{S}_{\leq r}(n) = \{X \in \text{Sym}(n) \mid \text{rank}(X) \leq r\}$$

Low-rank symmetric matrices

$$\mathcal{S}_{\leq r}(n) = \{X \in \text{Sym}(n) \mid \text{rank}(X) \leq r\} = \boxed{?}$$

- A decomposition

$$\mathcal{S}_{\leq r}(n) = \bigcup_{j=1}^{r+1} \mathcal{S}_j, \text{ with } \mathcal{S}_j := \{X \in \text{Sym}(n) \mid \lambda_j(X) = 0, \lambda_{j+n-r-1}(X) = 0\}$$

- Verify the error bound condition of $(\lambda_j, \lambda_{j+n-r-1})$
- Derivation of tangent sets $T_{\bigcup_{j=1}^{r+1} \mathcal{S}_j}(X) = \bigcup_{j=1}^{r+1} T_{\mathcal{S}_j}(X)$

Low-rank symmetric matrices

$$\mathcal{S}_{\leq r}(n) = \{X \in \text{Sym}(n) \mid \text{rank}(X) \leq r\} = \boxed{?}$$

- A decomposition

$$\mathcal{S}_{\leq r}(n) = \bigcup_{j=1}^{r+1} \mathcal{S}_j, \text{ with } \mathcal{S}_j := \{X \in \text{Sym}(n) \mid \lambda_j(X) = 0, \lambda_{j+n-r-1}(X) = 0\}$$

- Verify the error bound condition of $(\lambda_j, \lambda_{j+n-r-1})$
- Derivation of tangent sets $T_{\bigcup_{j=1}^{r+1} \mathcal{S}_j}(X) = \bigcup_{j=1}^{r+1} T_{\mathcal{S}_j}(X)$

Low-rank PSD matrices

$$\mathcal{S}_{\leq r}^+(n) = \{X \in \text{Sym}(n) \mid \lambda_{r+1}(X) = 0, \lambda_n(X) = 0\}$$

Low-rank symmetric matrices

$$\mathcal{S}_{\leq r}(n) = \{X \in \text{Sym}(n) \mid \text{rank}(X) \leq r\} = \boxed{?}$$

- A decomposition

$$\mathcal{S}_{\leq r}(n) = \bigcup_{j=1}^{r+1} \mathcal{S}_j, \text{ with } \mathcal{S}_j := \{X \in \text{Sym}(n) \mid \lambda_j(X) = 0, \lambda_{j+n-r-1}(X) = 0\}$$

- Verify the error bound condition of $(\lambda_j, \lambda_{j+n-r-1})$
- Derivation of tangent sets $T_{\bigcup_{j=1}^{r+1} \mathcal{S}_j}(X) = \bigcup_{j=1}^{r+1} T_{\mathcal{S}_j}(X)$

Low-rank PSD matrices


$$\mathcal{S}_{\leq r}^+(n) = \{X \in \text{Sym}(n) \mid \lambda_{r+1}(X) = 0, \lambda_n(X) = 0\}$$

$\mathcal{S}_{\leq r}^+(n)$ coincides with $\mathcal{S}_{r+1}$

Directional derivatives of singular values

Necessary notation [Zhang-Zhang-Xiao'13 SVVA]

- the SVD $X = \bar{U}[\bar{\Sigma}(X) \ 0] \bar{V}^\top$ with $\sigma_1(X) \geq \sigma_2(X) \geq \dots \geq \sigma_m(X)$
 $\alpha = \{i : \sigma_i(X) > 0, 1 \leq i \leq m\}$, $\beta = \{i : \sigma_i(X) = 0, 1 \leq i \leq m\}$
 $\beta_0 = \{m+1, \dots, n\}$
- the distinct singular values $\mu_1 > \mu_2 > \dots > \mu_p > \mu_{p+1} = 0$
 $\alpha_k = \{i : \sigma_i(X) = \mu_k, 1 \leq i \leq m\}$, $k = 1, \dots, p$, and $\hat{\beta} = \beta \cup \beta_0$



Directional derivatives of singular values (cont'd)

Four mappings


$$q_a : \{1, \dots, m\} \rightarrow \{1, \dots, p+1\}, q_a(i) = k, \text{ if } i \in \alpha_k$$

$$l : \{1, \dots, m\} \rightarrow \mathbb{N}, l(i) = i - \kappa_{q_a(i)-1}$$

$$q_b : \{1, \dots, m\} \rightarrow \mathbb{N}, q_b(i) = e, \text{ if } l(i) \in \beta_e^{q_a(i)}$$

$$l' : \{1, \dots, m\} \rightarrow \mathbb{N}, l'(i) = l(i) - \kappa_{q_b(i)-1}^{(q_a(i))}$$

Formula of $\sigma'_i(X; \eta)$ and $\sigma''_i(X; \eta, \zeta)$ at $X = \bar{U}[\bar{\Sigma}(X) \ 0] \bar{V}^\top$ for $i = 1, 2, \dots, m$



Compute second-order tangent set

Important geometry of $\mathcal{M}_{\leq r}$


- consider $X = \bar{U}[\Sigma \ 0] \bar{V}^\top \in \mathcal{M}_s$ with $\bar{U} = [U \ U_\perp]$ and $\bar{V} = [V \ V_\perp]$
- $\eta \in \text{T}_{\mathcal{M}_{\leq r}}(X)$ if and only if

$$\eta = U A V^\top + U_\perp B V^\top + U C V_\perp^\top + U_\perp R V_\perp^\top$$

where $\text{rank}(R) = \ell - s \leq r - s$

Index mappings associated with $\sigma_{r+1}(X) = 0$

- $r+1 \in \beta$, $q_a(r+1) = p+1$, $l(r+1) = r+1-s$
- $\bar{\eta}_{\beta\hat{\beta}} = U_\perp^\top \eta V_\perp = R$ and $\text{rank}(R) = \ell - s \Rightarrow q_b(r+1) = N_{p+1} + 1$
- $l'(r+1) = \ell + 1 - s$



Theorem

Given $X \in \mathcal{M}_{\leq r}$ and $\eta \in \mathbf{T}_{\mathcal{M}_{\leq r}}(X)$ with $\text{rank}(X) = s$ and the SVD $X = U \Sigma V^\top$. Let $\mathcal{P}_{\mathbf{N}_{\mathcal{M}_s}(X)}(\eta) = U_\eta \Sigma_\eta V_\eta^\top$, with $\text{rank}(\Sigma_\eta) = \ell - s$. Take $[U \ U_\eta \ U_{\eta^\perp}] \in \mathcal{O}(m)$ and $[V \ V_\eta \ V_{\eta^\perp}] \in \mathcal{O}(n)$. It holds that

$$\mathbf{T}_{\mathcal{M}_{\leq r}}^2(X; \eta) = \left\{ 2\eta X^\dagger \eta + [U^+ \ U_{\eta^\perp}] \begin{bmatrix} W_1 & W_2 \\ W_3 & J \end{bmatrix} [V^+ \ V_{\eta^\perp}]^\top \left| \begin{array}{l} W_1 \in \mathbb{R}^{\ell \times \ell} \\ W_2 \in \mathbb{R}^{\ell \times (n-\ell)} \\ W_3 \in \mathbb{R}^{(m-\ell) \times \ell} \\ J \in \mathbb{R}^{(m-\ell) \times (n-\ell)} \\ \text{rank}(J) \leq r - \ell \end{array} \right. \right\}$$

where $U^+ = [U \ U_\eta]$, $V^+ = [V \ V_\eta]$

A unified analysis of tangent sets to low-rank sets

Set	Format	First-order	Second-order
$\mathcal{M}$	general	Theorem 1	Theorem 1
$\mathcal{M}_{\leq r}$	matrix	[Schneider-USchmajew'15 SIOPT]	Proposition 1
$\mathcal{M}_{\leq r}^{\text{ht}}$	hierarchical Tucker	Proposition 2	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tc}}$	Tucker	[Gao-Peng-Yuan'25 MP]	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tt}}$	tensor train	[Kutschan'18 LAA]	Proposition 2
$\mathcal{S}_{\leq r}(n)$	symmetric matrix	[Li-Xiu-Zhou'20 JOTA]	Proposition 3
$\mathcal{S}_{\leq r}^+(n)$	PSD matrix	[Levin-Kileel-Boumal'25 MP]	Proposition 4

A unified analysis of tangent sets to low-rank sets

Set	Format	First-order	Second-order
$\mathcal{M}$	general	Theorem 1	Theorem 1
$\mathcal{M}_{\leq r}$	matrix	[Schneider-USchmajew'15 SIOPT]	Proposition 1
$\mathcal{M}_{\leq r}^{\text{ht}}$	hierarchical Tucker	Proposition 2	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tc}}$	Tucker	[Gao-Peng-Yuan'25 MP]	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tt}}$	tensor train	[Kutschan'18 LAA]	Proposition 2
$\mathcal{S}_{\leq r}(n)$	symmetric matrix	[Li-Xiu-Zhou'20 JOTA]	Proposition 3
$\mathcal{S}_{\leq r}^+(n)$	PSD matrix	[Levin-Kileel-Boumal'25 MP]	Proposition 4

Extension to intersection of sets $\mathcal{M} \cap \mathcal{K}$?

Theorem

- $\mathcal{M} = \{\tilde{X} \in \mathbb{R}^q \mid c(\tilde{X}) = 0\}$ with $c : \mathbb{R}^q \rightarrow \mathbb{R}^{n_1}$
- $\mathcal{K} = \{\tilde{X} \in \mathbb{R}^q \mid h(\tilde{X}) = 0\}$ with smooth $h : \mathbb{R}^q \rightarrow \mathbb{R}^k$

Theorem

- $\mathcal{M} = \{\tilde{X} \in \mathbb{R}^q \mid c(\tilde{X}) = 0\}$ with $c : \mathbb{R}^q \rightarrow \mathbb{R}^{n_1}$
- $\mathcal{K} = \{\tilde{X} \in \mathbb{R}^q \mid h(\tilde{X}) = 0\}$ with smooth $h : \mathbb{R}^q \rightarrow \mathbb{R}^k$
- Suppose that $(\mathcal{M}, \mathcal{K})$ satisfy

$$\begin{array}{ccc} \text{(transversal)} \quad \overline{\mathcal{M}} \cap \overline{\mathcal{K}} \subseteq \mathbb{R}^p & & \\ \text{(open)} \quad \phi \downarrow & \searrow h \circ \phi \text{ (rank-constant)} & \\ \mathcal{M} \cap \mathcal{K} \subseteq \mathbb{R}^q & \xrightarrow{h} & \mathbb{R}^k \end{array}$$

Theorem

- $\mathcal{M} = \{\tilde{X} \in \mathbb{R}^q \mid c(\tilde{X}) = 0\}$ with $c : \mathbb{R}^q \rightarrow \mathbb{R}^{n_1}$
- $\mathcal{K} = \{\tilde{X} \in \mathbb{R}^q \mid h(\tilde{X}) = 0\}$ with smooth $h : \mathbb{R}^q \rightarrow \mathbb{R}^k$
- Suppose that $(\mathcal{M}, \mathcal{K})$ satisfy

$$\begin{array}{ccc} \text{(transversal)} \quad \overline{\mathcal{M}} \cap \overline{\mathcal{K}} \subseteq \mathbb{R}^p & & \\ \text{(open)} \quad \phi \downarrow & \searrow h \circ \phi \text{ (rank-constant)} & \\ \mathcal{M} \cap \mathcal{K} \subseteq \mathbb{R}^q & \xrightarrow{h} & \mathbb{R}^k \end{array}$$

We have the following **intersection rules** for the tangent sets

$$\mathbf{T}_{\mathcal{M} \cap \mathcal{K}}(X) = \mathbf{T}_{\mathcal{M}}(X) \cap \mathbf{T}_{\mathcal{K}}(X)$$

$$\mathbf{T}_{\mathcal{M} \cap \mathcal{K}}^2(X; \eta) = \mathbf{T}_{\mathcal{M}}^2(X; \eta) \cap \mathbf{T}_{\mathcal{K}}^2(X; \eta)$$


Low-rank matrix sets $\mathcal{M}_{\leq r} \cap \mathcal{H}$

- $\mathcal{H} = \text{Aff}(m, n) = \{X \in \mathbb{R}^{m \times n} \mid \mathcal{A}(X) - b = 0\}$ [Li-Luo'23 SIOPT]
- $\mathcal{H} = \text{S}_F(m, n) = \{X \in \mathbb{R}^{m \times n} \mid \|X\|_F^2 - 1 = 0\}$ [Cason-Absil-Van Dooren'13 LAA]
- $\mathcal{H} = \text{Ob}(m, n) = \{X \in \mathbb{R}^{m \times n} \mid \text{diag}(XX^\top) - \mathbf{1} = \mathbf{0}\}$ [Yang-Gao-Yuan'25]
- $\mathcal{H} = \mathbb{H}_{n-1}^m = \{X \in \mathbb{R}^{m \times n} \mid \text{diag}(X \text{Diag}(-1, 1, \dots, 1)X^\top) + \mathbf{1} = \mathbf{0}\}$ ☺ new

A smooth parameterization

$$(\mathcal{M}_{\text{LR}}, \phi_{\text{LR}}) = (\mathbb{R}^{m \times r} \times \mathbb{R}^{n \times r}, (L, R) \mapsto LR^\top)$$

- ☺ $\mathcal{M}_{\text{LR}}$ coincides with the ambient space
- ☺ ϕ_{LR} is open around “balanced” (L, R)
- ★ It remains to prove $\overline{\mathcal{H}} = \phi_{\text{LR}}^{-1}(\mathcal{H})$ is manifold



A unified analysis of tangent sets

Set	Format	First-order	Second-order
$\mathcal{M}$	Assumption 1	Theorem 1	Theorem 1
$\mathcal{M}_{\leq r}$	matrix	[Schneider-USchmajew'15 SIOPT]	Proposition 1
$\mathcal{M}_{\leq r}^{\text{ht}}$	hierarchical Tucker	Proposition 2	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tc}}$	Tucker	[Gao-Peng-Yuan'25 MP]	Proposition 2
$\mathcal{M}_{\leq r}^{\text{tt}}$	tensor train	[Kutschan'18 LAA]	Proposition 2
$\mathcal{S}_{\leq r}(n)$	symmetric matrix	[Li-Xiu-Zhou'20 JOTA]	Proposition 3
$\mathcal{S}_{\leq r}^+(n)$	PSD matrix	[Levin-Kileel-Boumal'25 MP]	Proposition 4
Intersection of sets	Structured set	First-order	Second-order
$\mathcal{M} \cap \mathcal{K}$	Assumption 2	Theorem 2	Theorem 2
$\mathcal{M}_{\leq r} \cap \mathcal{H}$	$\mathcal{H}$ is an affine manifold	[Li-Luo'23 SIOPT]	Appendix C.1
$\mathcal{M}_{\leq r} \cap \mathcal{H}$	$\mathcal{H}$ is orthogonally invariant	[Yang-Gao-Yuan'25]	Appendix C.2
$\mathcal{M}_{\leq r} \cap \mathcal{H}$	$\mathcal{H}$ is hyperbolic	Appendix C.3	Appendix C.3
$\mathcal{S}_{\leq r}(n) \cap \mathcal{U}$	$\mathcal{U} = \{X \mid \ X\ _{\text{F}}^2 = 1\}$	Appendix D.2	Appendix D.2
$\mathcal{S}_{\leq r}^+(n) \cap \mathcal{U}$	$\mathcal{U} = \{X \mid \mathcal{A}(X) = b\}$	[Levin-Kileel-Boumal'25 MP]	Appendix D.3


Tangent sets bridge optimization landscapes

Second-order optimality conditions

Second-order stationarity [Ruszczynski'06]

- $-\nabla f(X^*) \in \hat{N}_{\mathcal{M}}(X^*)$
- $\langle \nabla f(X^*), \zeta \rangle + \langle \eta, \nabla^2 f(X^*)[\eta] \rangle \geq 0$, for every $\eta \in T_{\mathcal{M}}(X)$ such that $\langle \nabla f(X^*), \eta \rangle = 0$ and all $\zeta \in T_{\mathcal{M}}^2(X^*; \eta)$

Smooth parameterization finds first-order stationary points [Levin-Kileel-Boumal'25 MP]



$$\min_{Y \in \overline{\mathcal{M}}} \bar{f}(Y) = f(\phi(Y))$$

$$\min_{X \in \mathcal{M}} f(X)$$

When does a parameterization satisfy “ $2 \Rightarrow 2$ ”?

A sufficient and necessary condition for “ $2 \Rightarrow 2$ ”

Mappings connecting the geometry

$$\mathbf{L}_Y : T_{\overline{\mathcal{M}}}(Y) \rightarrow \mathcal{E} : v \mapsto D\phi_Y[v], \text{ for } Y \in \overline{\mathcal{M}}$$

$$\mathbf{Q}_{Y,v} : T_{\overline{\mathcal{M}}}^2(Y; v) \rightarrow \mathcal{E} : u_v \mapsto D\phi_Y[u_v] + D^2\phi_Y[v, v], \text{ for } Y \in \overline{\mathcal{M}} \text{ and } v \in T_{\overline{\mathcal{M}}}(Y)$$

Two important sets

$$\mathcal{Q}_Y(v) := \bigcup_{\{v_i\}_{i \in \mathbb{N}} : \mathbf{L}_Y(v_i) \rightarrow \mathbf{L}_Y(v)} \lim_{i \rightarrow \infty} (\mathbf{Q}_{Y,v_i}(u_{v_i}) + \text{im}(\mathbf{L}_Y)), \text{ for } v \in T_{\overline{\mathcal{M}}}(Y)$$

$$\Gamma_Y := \left\{ \mathcal{P}_{\hat{\mathbf{N}}_{\mathcal{M}}(\phi(Y))} \left(D\phi_Y[\Pi(v_0, v_1)] + D^2\phi_Y[v_0, v_1] \right) \mid v_0 \in \ker(\mathbf{L}_Y), v_1 \in T_{\overline{\mathcal{M}}}(Y) \right\}$$

Theorem

The parameterization $(\overline{\mathcal{M}}, \phi)$ satisfies “ $2 \Rightarrow 2$ ” at $Y \in \overline{\mathcal{M}}$ if and only if $\text{im}(\mathbf{L}_Y) = T_{\mathcal{M}}(X)$ where $X = \phi(Y)$, and for all $v \in T_{\overline{\mathcal{M}}}(Y)$

$$T_{\mathcal{M}}^2(X; \mathbf{L}_Y(v)) \subseteq \overline{\text{conv}}(\mathcal{Q}_Y(v) + \mathcal{Q}_Y(0) + \Gamma_Y)$$

Second-order optimality on bounded-rank matrices

Proposition

$$\begin{array}{ll} \min_{X \in \mathbb{R}^{m \times n}} & f(X) \\ \text{s. t.} & X \in \mathcal{M}_{\leq r} \end{array}$$

The point $X^* \in \mathcal{M}_{\leq r}$ with $\text{rank}(X^*) = s$ is **second-order stationary** if

$$\begin{cases} \nabla_{\mathcal{M}_r} f(X^*) = 0 \text{ and } \nabla_{\mathcal{M}_r}^2 f(X^*) \succeq 0, & \text{if } s = r \\ \nabla f(X^*) = 0 \text{ and } \langle \eta, \nabla^2 f(X^*)[\eta] \rangle \geq 0 \text{ for all } \eta \in \text{T}_{\mathcal{M}_{\leq r}}(X^*), & \text{if } s < r \end{cases}$$

Complexity of verifying second-order conditions (SOC)

Find a negative curvature direction at X with $\text{rank}(X) < r$

$$\begin{aligned} \min_{\eta \in \mathbb{R}^{m \times n}} \quad & \langle \eta, \mathcal{A}(\eta) \rangle \\ \text{s. t.} \quad & \|\eta\|_F = 1 \\ & \eta \in \mathcal{T}_{\mathcal{M}_{\leq r}}(X) \end{aligned} \tag{Q}$$

Problem: VERSOC

Input: Parameters m, n, r ; point $X \in \mathcal{M}_{\leq r}$; symmetric operator $\mathcal{A}$

Question: Does the optimal value of (Q) $\lambda^* < 0$?

Complexity of verifying second-order conditions (SOC)

Find a negative curvature direction at X with $\text{rank}(X) < r$

$$\begin{aligned} \min_{\eta \in \mathbb{R}^{m \times n}} \quad & \langle \eta, \mathcal{A}(\eta) \rangle \\ \text{s. t.} \quad & \|\eta\|_F = 1 \\ & \eta \in \mathcal{T}_{\mathcal{M}_{\leq r}}(X) \end{aligned} \tag{Q}$$

Problem: VERSOC

Input: Parameters m, n, r ; point $X \in \mathcal{M}_{\leq r}$; symmetric operator $\mathcal{A}$

Question: Does the optimal value of (Q) $\lambda^* < 0$?

An **NP-complete** problem [Karp'72]

Problem: CLIQUE

Input: Undirected graph $G = (\mathcal{V}, E)$; clique size K

Question: Does there exist a clique of size K in G ?

Theorem (NP-hardness)

The problem CLIQUE is polynomially reducible to VERSOC, and thus verifying second-order optimality conditions is **NP-hard**

Theorem (NP-hardness)

The problem CLIQUE is polynomially reducible to VERSOC, and thus verifying second-order optimality conditions is **NP-hard**

Theorem (No FPTAS)

Unless $P=NP$, there is **no fully polynomial-time approximation scheme** for verifying whether a point is second-order stationary for optimization problems over bounded-rank matrices

A glimpse: optimization over the intersection of manifolds

Problem formulation


$$\begin{array}{ll} \min_{X \in \mathcal{E}} & f(X) \\ \text{s. t.} & X \in \mathcal{H} \cap \mathcal{M} \end{array}$$

Theory

Intrinsic transversality is equivalent to clean intersection

$$\mathbf{T}_{\mathcal{H} \cap \mathcal{M}}(X) = \mathbf{T}_{\mathcal{H}}(X) \cap \mathbf{T}_{\mathcal{M}}(X)$$

Geometric method



Take-home message

- ★ Unified analysis for tangent sets
- ★ Second-order optimality conditions
- ★ NP-hardness of achieving second-order stationarity
- ★ Bridges between optimization landscapes based on tangent sets

Perspective

- 😊 Efficient algorithm exploiting the geometry [Yang-Gao-Yuan'26]

$$\mathbf{T}_{\mathcal{H} \cap \mathcal{M}}(X) = \mathbf{T}_{\mathcal{H}}(X) \cap \mathbf{T}_{\mathcal{M}}(X)$$

- 🔗 Second-order development beneficial from the property

$$\mathbf{T}_{\mathcal{H} \cap \mathcal{M}}^2(X; \eta) = \mathbf{T}_{\mathcal{H}}^2(X; \eta) \cap \mathbf{T}_{\mathcal{M}}^2(X; \eta)$$

Thanks for your attention!

✉ yangyan1801@gmail.com

👤 <https://ucas-yanyang.github.io/>

📅 Looking for postdoc positions from July, 2027



Scan for homepage

References

Yan Yang, Bin Gao, Ya-xiang Yuan

- [1] Variational analysis of determinantal varieties. arXiv:2511.22613 2025
- [2] *A space-decoupling framework for optimization on bounded-rank matrices with orthogonally invariant constraints*. Mathematical Programming (2026)
- [3] Optimization over the intersection of manifolds. arXiv:2605.22736 2026
- [4] Bilevel reinforcement learning via the development of hyper-gradient without lower-level convexity. AISTATS (2025)
- [5] LancBiO: dynamic Lanczos-aided bilevel optimization via Krylov subspace. ICLR (2025)